

FIRST RECORD OF EDIBLE LARVAE OF RHINOSTOMUS BARBIROSTRIS (FABRICIUS, 1775) (COLEOPTERA: DRYOPHTHORIDAE) ON OENOCARPUS BATAUA MART. IN THE STATE OF ACRE, BRAZIL

PRIMEIRO REGISTRO DE LARVAS COMESTÍVEIS DE RHINOSTOMUS BARBIROSTRIS (FABRICIUS, 1775) (COLEOPTERA: DRYOPHTHORIDAE) EM OENOCARPUS BATAUA MART. NO ESTADO DO ACRE, BRASIL

PRIMER REGISTRO DE LARVAS COMESTIBLES DE RHINOSTOMUS BARBIROSTRIS (FABRICIUS, 1775) (COLEOPTERA: DRYOPHTHORIDAE) EN OENOCARPUS BATAUA MART. EN EL ESTADO DE ACRE, BRASIL



<https://doi.org/10.56238/arev7n11-395>

Submission date: 10/30/2025

Publication Date: 11/30/2025

Dagmar Mercado Soares¹, Evandro José Linhares Ferreira², Fernanda Grazielly Soares Lima³, Adalberto Hipólito de Sousa⁴, Guaracy Barbosa dos Santos Maia⁵, William Ferreira Alves⁶, Eraldo Medeiros Costa Neto⁷, Anselmo Fortunato Ruiz Rodriguez⁸

ABSTRACT

This study presents the first record of edible larvae of *Rhinostomus barbirostris* on the *Oenocarpus bataua* palm tree in Acre, Brazil. The record was made in the native forest of the Chico Mendes Extractive Reserve (RESEX) in the municipality of Epitaciolândia. The larval record expands knowledge about the distribution of the species in the southwestern Amazonia and represents the first step toward future research focused on their bioprospecting.

Keywords: Edible Insects. Coleoptera. Arecaceae. Southwestern Amazon.

RESUMO

Este estudo apresenta o primeiro registro da ocorrência de larvas comestíveis de *Rhinostomus barbirostris* na palmeira *Oenocarpus bataua* no Acre, Brasil. O registro foi realizado em floresta nativa da Reserva Extrativista Chico Mendes (RESEX) situada no município de Epitaciolândia. O registro das larvas expande o conhecimento sobre a

¹ Doctoral student in Biotechnology. Universidade Federal do Acre. Acre, Brazil.

E-mail: dagmamercado@gmail.com

² Dr. of Botany. Instituto Nacional de Pesquisas da Amazônia (INPA). Acre, Brazil.

³ Specialization in Biotechnology. Centro Universitário Uninorte. Acre, Brazil.

E-mail: fernandagraziellyac@gmail.com

⁴ Dr. of Entomology. Universidade Federal do Acre. Acre, Brazil.

E-mail: adalberto.sousa@ufac.br

⁵ Master's degree in Chemical Engineering. Universidade Federal do Acre. Acre, Brazil.

E-mail: guaracy.maia@ufac.br

⁶ Dr. of Chemistry. Universidade Federal do Acre. Acre, Brazil.

E-mail: william.alves@ufac.br

⁷ Dr. of Ecology. Universidade Estadual de Feira de Santana. Bahia, Brazil.

E-mail: eraldont@uefs.br

⁸ Postdoctoral Researcher in Nanoscience and Nanotechnology. Universidade Federal do Acre. Acre, Brazil.

E-mail: anselmo.rodriguez@ufac.br

distribuição da espécie no sudoeste da Amazônia e representa a primeira etapa para futuras pesquisas voltadas para a bioprospecção das mesmas.

Palavras-chave: Insetos Comestíveis. Coleoptera. Arecaceae. Sudoeste da Amazônia.

RESUMEN

Este estudio presenta el primer registro de larvas comestibles de *Rhinostomus barbirostris* en la palmera *Oenocarpus bataua* en Acre, Brasil. El registro se realizó en el bosque nativo de la Reserva Extractivista Chico Mendes (RESEX), en el municipio de Epitaciolândia. El registro de las larvas amplía el conocimiento sobre la distribución de la especie en el suroeste de la Amazonía y representa el primer paso para futuras investigaciones orientadas a su bioprospección.

Palabras clave: Insectos Comestibles. Coleoptera. Arecaceae. Suroeste de la Amazonía.

1 INTRODUCTION

Rhinostomus barbirostris (Fabricius, 1775) (Coleoptera: Dryophthoridae), is a holometabolous insect with a neotropical distribution that generally coincides with the palm family (Arecaceae) (Morrone; Cuevas, 2002; Delgado *et al.*, 2008; Casas-Reátegui *et al.*, 2018). Although adults are commonly documented (Setz, 1991; Choo *et al.*, 2009; Araujo *et al.*, 2018), immature forms, especially the larval stage, remain poorly understood, particularly in the Brazilian Amazon (Coimbra Junior, 1984; Molina *et al.*, 2023).

Larvae of *R. barbirostris* can develop in palm tree stem of Arecaceae species such as *Oenocarpus bacaba*, *Mauritia flexuosa*, *Oenocarpus bataua*, *Attalea phalerata*, among others, after natural or intentional fall (Coimbra Junior, 1984; Choo *et al.*, 2009; Casas-Reátegui *et al.*, 2018). In some regions of the Amazon, the larvae are considered a food resource by traditional peoples and communities and can be eaten raw, boiled, roasted, or fried in their own oil (Coimbra Junior, 1984; Araujo; Beserra, 2007; Choo *et al.*, 2009; Casas-Reátegui *et al.*, 2018; Cerna *et al.*, 2021; Jaramillo-Vivanco *et al.*, 2023).

Considering that there were no official records of the occurrence of *R. barbirostris* beetle larvae in the state of Acre, in the southwestern Brazilian Amazon, this study aims to present the first record of these larvae found in the stems of *O. bataua* palm trees growing spontaneously in local native forests.

2 MATERIAL AND METHODS

The larvae were found and collected from the stems of adult *O. bataua* palm trees growing spontaneously in a native forest area located in the rubber tapper settlement called “Pinheiro”, located near the “Riozinho” stream (10°44'19.4“S; 68°48'56.1“W), within the Chico Mendes Extractive Reserve (RESEX), in the municipality of Epitaciolândia, Acre, Brazil (Figure 1).

Figure 1

Location of the larvae collection point



Sources: SIRGAS Datum (2000); IBGE (2022); Google Earth Engine (2025).

The collection of biological material was authorized and carried out as part of research registered with the Biodiversity Authorization and Information System (SISBIO 87534-1 and 87970-1) and the National System for the Management of Genetic Heritage and Associated Traditional Knowledge (SisGen A33BAE0). The specimen of *O. bataua* botanical material was deposited in the Herbarium of the Zoobotanical Park (UFACPZ), under number 30555.

To collect the larvae, it was necessary to fell an adult *O. bataua* palm tree. On the ground, using an axe, the trunk was cut crosswise into smaller pieces, which were then split lengthwise to facilitate the collection of as many intact larvae as possible (Figura 2A–F).

3 RESULTS AND DISCUSSION

In the middle region of the palm tree stem, galleries containing numerous live larvae of *R. barbirostris* were observed (Figure 2A–C). These larvae were cream-colored, soft, and greasy in consistency, with a characteristic odor (Figure 2D–F).

Figure 2

A *Oenocarpus bataua* stem being opened with the help of an axe; *B* - Numerous *Rhinostomus barbirostris* larvae in the stem galleries; *C* - Close up of a larva in a sample gallery of the stem; *D* - Larva collected; *E* and *F* - Larvae collected and packed in a container.




Source: Prepared by the authors.

At the same time, numerous pupae and some adult forms were found in the stem apical region, showing that the entire development cycle of the Dryophthoridae occurs inside the stem of this palm tree (Figure 3A–F).

Figure 3

A - Pupae of *Rhinostomus barbirostris* in the stem galleries; B - Pupae in the stem; C - Pupae collected; D - Adults in the stem; E - Adults collected; F - Adults of *R. barbirostris*, duly identified, female (left) and male (right)



Source: Prepared by the authors.

Rhinostomus barbirostris (Fabricius, 1775) (Coleoptera: Dryophthoridae). New Record: BRAZIL. Acre: Epitaciolândia, Chico Mendes Extractive Reserve (RESEX), “Colocação” Pinheiro, near Igarapé Riozinho, terra firme forest (10°44'19.4”S and 68°48'56.1”W), 250 m altitude, June 20, 2024 (pupa and adult), D.M. Soares (insect collection number: 01) (UFACLE.2024.01).

In the state of Rondônia, in the southwestern Brazilian Amazon, the same larvae were recorded on the trunk of *O. bataua* (synonym: *Jessenia bataua*), where they are locally called “mãyõrã” by the Paiter Suruí indigenous people (Coimbra Junior, 1984).

Although documented in other regions of the Amazon region (Choo *et al.*, 2009), there was still no record of the presence of these larvae in association with palm trees in the state of Acre, Brazil. Thus, documenting the presence of immature forms of *R. barbirostris* on the trunk of the *O. bataua* palm tree filled a gap in knowledge about its occurrence in the state of Acre, since until now only adult forms of this species had been recorded in Acre (Machado; Liesenfeld, 2019).

4 CONCLUSION

The photographic images of *R. barbirostris* larvae consolidate the first official document of its occurrence in the state of Acre and a new record for the southwestern Brazilian Amazon. This discovery constitutes the initial stage for future research focused on their bioprospecting.

ACKNOWLEDGEMENTS

The first author would like to thank the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), the Graduate Program in Biodiversity and Biotechnology of the Amazon, Rede Bionorte, and Mr. Rildo Coelho de Figueiredo for access to biological material.

REFERENCES

- Araujo, J. J., Keller, H. A., & Hilgert, N. I. (2018). Management of pindo palm (*Syagrus romanzoffiana*, Arecaceae) in rearing of Coleoptera edible larvae by the Guarani of Northeastern Argentina. *Ethnobiology and Conservation*, 7(1), 1–18.
- Araujo, Y., & Beserra, P. (2007). Diversidad en invertebrados consumidos por las etnias Yanomami y Yekuana del Alto Orinico, Venezuela. *Interciencia*, 32(5), 318–323.
- Casas-Reátegui, R., et al. (2018). Beetles, ants, wasps, or flies? An ethnobiological study of edible insects among the Awajún Amerindians in Amazonas, Peru. *Journal of Ethnobiology and Ethnomedicine*, 14(1), 1–11.
- Cerna, A., et al. (2021). Traditional knowledge regarding entomophagy in San Martín, Peruvian Amazon. *Peruvian Journal of Agronomy*, 5(2), 44–59.
- Choo, J., Zent, E. L., & Simpson, B. B. (2009). The importance of traditional ecological knowledge for palm-weevil cultivation in the Venezuelan Amazon. *Journal of Ethnobiology*, 29(1), 113–128.
- Coimbra Júnior, C. E. A. (1984). Estudos de ecologia humana entre os Suruí do Parque Indígena Aripuanã, Rondônia: 1. O uso de larvas de Coleópteros (Bruchidae e Curculionidae) na alimentação. *Revista Brasileira de Zoologia*, 2(2), 35–47.

- Delgado, C., Couturier, G., Mathews, P., & Mejia, K. (2008). Comercialización de larvas de *Rhynchophorus palmarum* (Coleoptera: Dryophthoridae) en la Amazonía peruana. *Boletín de la Sociedad Entomológica Aragonesa*, 42, 407–412.
- Jaramillo-Vivanco, T., Cámara, R. M., Cámara, M., Tejera, E., Balslev, H., & Álvarez-Suárez, J. M. (2023). Ethnobiology of edible palm weevil larvae *Rhynchophorus palmarum* L. (Curculionidae, Coleoptera), a common food source in Amazonian Ecuador. *Journal of Insects as Food and Feed*, 10(13), 427–441.
- Machado, E., & Liesenfeld, M. V. A. (2019). Registro da presença da broca-da-estipe-do-coqueiro *Rhynchophorus palmarum* e da broca-do-tronco *Rhinostomus barbirostris* (Curculionidae: Coleoptera) em Cruzeiro do Sul (Acre) e seu potencial de danos. In *Anais do 1.º Simpósio Acreano de Espécies Exóticas Invasoras*. Even3. <https://www.even3.com.br/anais/simposioexoticasinvasoras2019/171700-registro-da-presenca-da-broca-da-estipe-do-coqueiro-rhynchophorus-palmarum-e-da-broca-do-tronco-rhinostomus-barbi/>
- Molina, A. A., Ferreira, M. J., Suruí, L. O., Norder, L. A. C., Costa Neto, E. M., & Clement, C. R. (2023). Amazonian invertebrates in the traditional diet of the Paiteir Suruí in southeastern Brazil. *Human Ecology Interdisciplinary Journal*, 51(2), 209–219.
- Morrone, J. J., & Cuevas, P. (2002). Cladistics of the pantropical genus *Rhinostomus* (Coleoptera: Curculionoidea: Dryophthoridae), with nomenclatural notes. *Journal of the New York Entomological Society*, 110, 376–388.
- Setz, E. Z. F. (1991). Animals in the Nambiquara diet: Methods of collection and processing. *Journal of Ethnobiology*, 11(1), 1–22.
- Vaurie, P. (1970). Weevils of the tribe Sipalini (Coleoptera, Curculionidae, Rhynchophorinae). Part 1: The genera *Rhinostomus* and *Yuccaborus*. *American Museum Novitates*, (2419), 1–57.