


EMPIRICAL INDICATORS FOR THE CONSTRUCTION OF SOFTWARE FOR THE IMPLEMENTATION OF THE SYSTEMATIZATION OF NURSING CARE IN ASSISTED REPRODUCTION

 <https://doi.org/10.56238/arev6n2-159>

Submitted on: 17/09/2024

Publication date: 17/10/2024

Michelly Guedes de Oliveira Araújo¹, Rosilene Santos Baptista², Michelinne Oliveira Machado Dutra³, Ana Carolina Dantas Rocha Cerqueira⁴, Francisco Stélio de Sousa⁵, Adriana Gomes Nogueira Ferreira⁶, Carla Carolina da Silva Leite Freitas⁷ and Inácia Sátiro Xavier de França⁸

¹ Dr. in Nursing

HUAC

Email: michelly.g@gmail.com

Orcid: <https://orcid.org/0000-0002-2904-3878>

Lattes: <http://lattes.cnpq.br/6384337782089345>

² Dr. in Nursing

UEPB:

E-mail: rosilenesbaptista@gmail.com

Orcid: <https://orcid.org/0000-0001-7400-7967>

Lattes: <http://lattes.cnpq.br/9884386554804972>

³ Dr. in Nursing

UEPB

Email: michelinne_oliveira@hotmail.com

Orcid: <https://orcid.org/0000-0003-4066-8964>

Lattes: <http://lattes.cnpq.br/439763771561023>

⁴ Dr. in Nursing

UEPB

E-mail: ana.cerqueira@servidor.uepb.edu.br

Orcid: <https://orcid.org/0000-0001-5782-3102>

Lattes: <http://lattes.cnpq.br/4750070753228132>

⁵ Doctor of Nursing

UEPB

Email: stelio@servidor.uepb.edu.br

Orcid: <https://orcid.org/0000-0002-9604-4133>

Lattes: <http://lattes.cnpq.br/6010622299643929>

⁶ Dr. in Nursing

UFMA

E-mail: adriana.nogueira@ufma.br

Orcid: <https://orcid.org/0000-0002-7107-1151>

Lattes: <http://lattes.cnpq.br/4245835067415813>

⁷ Master in Development and Environment

UFPB

E-mail: carlacarolina01@gmail.com

Lattes: <http://lattes.cnpq.br/4180598396678659>

Orcid: <https://orcid.org/0000-0002-4780-7640>

⁸ Dr. in Nursing

UEPB:

E-mail: inacia.satiro@servidor.uepb.edu.br

Orcid: <https://orcid.org/0000-0002-2695-510X>

Lattes: <http://lattes.cnpq.br/2783760383527777>

ABSTRACT

Infertility is conceptualized by the inability to develop a successful pregnancy through sexual intercourse or artificial insemination in a period of 12 months or more. Nursing works during the process of implementing assisted reproduction treatments. In this sense, the use of software provides support in the documentation of the actions that govern care, in addition to assisting in the elaboration of clinical judgment, which can result in better quality care. Thus, the objective of this study was to identify the empirical indicators of women submitted to assisted reproduction techniques in order to structure a support software for the Systematization of Nursing Care for these women. This is a methodological research through the identification of empirical indicators related to women submitted to Assisted Reproduction techniques. As a result, a sample of 40 articles was obtained, from which 166 indicators were extracted. A total of 166 empirical indicators were obtained, of which 74 represented psychobiological needs, 88 psychosocial needs and 4 psychospiritual HEIs. Therefore, from the identification of the altered empirical indicators of women submitted to assisted reproduction techniques in the light of the theory of Basic Human Needs, it will be possible to develop a valid software for Systematization of Nursing Care to women submitted to assisted reproduction techniques.

Keywords: Assisted reproduction techniques, Nursing Process, Nursing, Software, Health technology.

INTRODUCTION

According to the American Society for Reproductive Medicine (ASRM), infertility is conceptualized as the inability to develop a successful pregnancy through sexual intercourse or artificial insemination within a period of 12 months or more (Practice Committee of the American Society for Reproductive Medicine, 2013).

In the global context, infertility remains a highly prevalent constant. According to the WHO (2011), it affects 8 to 15% of couples of reproductive age, and is characterized as a growing condition due to factors that delay motherhood and ultimately affect the fertility of spouses. It has an overall incidence of 9% on average, reaching rates of 50% in some regions of Central Africa, 11% in the developed world, and 20% in the Mediterranean region (Boivin et al., 2007; Jumayev et al., 2012).

When analyzing infertility from a macro context, it should be considered a relevant public health problem and inserted as a disease. And not to approach it in a simplistic way, seeing it only as a setback that affects family and individual well-being and the social insertion of affected couples (Cook; Dickens, 2014). Infertility can have harmful repercussions on women's emotional state, as well as on physical health (Mendonça et al., 2014).

In order to solve infertility by enabling infertile couples to become pregnant and to encompass homosexual relationships, reproductive technologies (RT) emerged in the 1980s. These individuals project, in the existing techniques, possibilities of forming a family with descendants, and thus having symbolic and social values that govern a family, even if they do not experience reproductive sexuality (Machin; Couto, 2014).

From this perspective, men and women refer to the development of assisted reproductive technologies as a strategy to compensate for the decrease in fertility due to advancing age, as it facilitates the procreation process. However, these individuals are unaware of the onerous costs of treatment and the proportion of success rates, as positive pregnancy results are not always achieved, they only demonstrate high confidence in the treatment that will be carried out (Daniluk; Koert, 2013; CFM, 2013).

It is possible to mention several activities performed by nurses in assisted reproduction treatments, even though they are not the professional who certifies the diagnosis and provides medication prescriptions (Queiroz et al., 2020). But it is the one who acts in welcoming, qualified listening to frustrated attempts to get pregnant; collects the necessary data from your gynecological and obstetric history; identifies the nursing

diagnoses applicable to each specificity, whether physical or psychological; plans interventions aimed at each change detected; outlines the goals and evaluates the evolution of this patient throughout the treatment. In addition, it provides clarification on the conducts that will be taken, administers medications, inspects their effects, and makes itself available to patients in order to enable quality health care (Queiroz et al., 2020).

In view of the scenario presented by infertility, imbalances permeate the individual in this condition and raise needs. Aiming at this context, it was sought to associate the concepts that excel in the balance of the human being and provide a complete well-being contained in the Theory of Basic Human Needs (NHB) by Wanda Horta.

This theory was created based on Maslow's Theory of Human Motivation, which is based on NHB. They are categorized into five levels: physiological needs, security, love, esteem, and self-actualization.

Thus, according to Horta (2015), an individual only seeks to meet the subsequent categories after being minimally satisfied in the previous ones. However, this satisfaction is never complete or lasting, otherwise there would be no individual motivation.

Regarding BHN, these are defined as levels of tension, conscious or unconscious, resulting from hemodynamic imbalances of vital phenomena. They are exacerbated when an established imbalance occurs, which may be apparent, conscious, verbalized or not by an individual, his family or the community (Horta, 2015).

Empirical indicators (IES) comprise the instruments of research practices or tools, used to assess concepts of a given theory (Fawcett, 2021). Thus, the study is pertinent due to the possibility of collaborating with the Systematization of Nursing Care for women submitted to assisted reproduction

In view of the above, the objective of this study was to identify the empirical indicators of women submitted to assisted reproduction, in order to structure a software for the Systematization of Nursing Care, in the light of the Theory of Basic Human Needs.

METHODOLOGY

This is a methodological study, developed through the survey of altered empirical indicators of women submitted to assisted reproduction. These were obtained through an integrative review of the scientific literature following the following specifications: identification of the problem, definition of the research question, establishment of inclusion and exclusion criteria, definition of the information to be extracted from the studies,

evaluation of the included studies, interpretation of the results and synthesis of data and presentation (Soares et al., 2014).

The methodological study is intended for investigations, organization and analysis of data, elaboration, validation and evaluation of research instruments and techniques, to enable the obtaining of a reliable, accurate and usable instrument so that it can be replicated by other researchers (Polit et al., 2019).

Thus, the following guiding question was defined for the integrative review: "what are the basic human needs of Wanda Horta's Theory found in women undergoing assisted reproduction techniques?"

The selection of articles was made through the following libraries and databases: Scientific Electronic Library Online (SciELO), Pubmed, Sciverse Scopus – Elsevier (Scopus), Latin American and Caribbean Health Sciences Information System (LILACs), Web of Science, Cochrane Library, Science Direct and Nursing Database (BDENF) via Virtual Health Library (VHL).

The search for studies in the bases took place between March and June 2020. It was performed in an uncontrolled manner through the descriptors identified in the Health Sciences Descriptors (DeCS) and in the Medical Subject Heading Terms (MeSH) through the following crosses: 1 (assisted reproductive techniques, nursing) / (assisted reproductive techniques, nursing) / (assisted reproductive techniques, nursing) and 2 (assisted reproductive techniques, women's health, nursing) / (assisted reproductive techniques, women's health, nursing) / (assisted reproductive techniques, women's health, nursing). English, Portuguese and Spanish languages were included.

The following inclusion criteria were adopted: articles available in full, which answered the guiding question, in Portuguese, English, and Spanish, between the years 2015 and 2020. And as exclusion criteria: repeated articles, uncompleted research, editorials, book chapters, expert opinions.

RESULTS AND DISCUSSION

Following the determined parameters, the first search obtained 1,305 studies, after applying the inclusion criteria resulted in 110, with the application of the exclusion criteria, 61 articles and the final sample totaled 40 articles after reading in full, as shown in table 1.

Table 1. Studies available in the databases acquired from the integrative literature review.

Databases	Studies acquired with crosses	Studies after application of the inclusion criteria	Studies after application of exclusion criteria	Studies after reading in full
Scielo	1	0	0	0
Pubmed	173	13	11	11
Scopus	657	57	34	22
Lilacs	6	3	0	0
Web of Science	32	7	1	1
Cochrane	225	4	3	0
Science Direct	211	26	12	6
BDENF	0	0	0	0
TOTAL	1.305	110	61	40

Source: Prepared by the authors.

Thus, the objective of this study was to recognize the altered manifestations of NHB in women undergoing assisted reproduction through an integrative literature review. The final sample totaled 40 articles. Of these, 11 were from Pubmed, 22 from Scopus, 1 from Web of Science and 6 from Science Direct.

Studies were selected between 2015 and 2020, with a predominance in 2019 with 15 publications (37.5%), followed by 2017 and 2018 with 5 studies in each year (12.5%), 9 in 2016 (22.5%), 4 in 2015 (10%) and 2 in 2020 (5%). It was also found that 90% of the articles were international studies. Regarding the methodological design, quantitative studies were the predominant ones with 35%, followed by qualitative studies with 15%, as shown in Table 2 below.

Table 2. Distribution of the types of methodological designs and their frequency according to the integrative literature review.

Methodological design	N	%
Quantitative	14	35
Qualitative	6	15
Quantum-Which	1	2,5
Revision	5	12,5
Systematic review	3	12,5
Systematic review and meta-analysis	3	12,5
Randomized controlled trial	3	7,5
Almost experimental	1	2,5

Source: Prepared by the authors.

To emphasize the conceptualization of empirical indicators, Denser (2003) states that in contexts within normality or in conditions within the expected, indicators expose the modifications, and it is from their detection that one can act in the prevention of problems. Thus, this study identified inadequacies of the basic human needs of the public that

undergoes assisted reproduction treatments so that it was possible to act emphatically on these points and minimize the damage caused.

Dealing with the levels of human needs, psychobiological and psychosocial needs are shared by all individuals, but psychospiritual needs are unique characteristics of the human being (Neto et al., 2019).

Thus, operational definitions were developed based on the analysis of studies in the integrative literature review. The empirical indicators were distributed in the classifications of the Theory of Wanda Horta (2015) adjusted by Garcia and Cubas (2012) and captured by the identification of these altered and subdivided NBN by the levels.

A total of 166 empirical indicators (HEIs) were obtained, of which 74 represented psychobiological needs, 88 altered psychosocial needs, and 4 psychospiritual HEIs, as shown in tables 3, 4, and 5 below.

In this first level of the theory addressed, some needs were not contemplated with indicators, as it was not detected in the literature altered human needs of: oxygenation, hydration, body care, physical integrity and thermal regulation.

Thus, with regard to nutrition, some changes were pointed out, such as: obesity, inadequate nutrition, and changes in appetite. The studies by Moura, Souza & Scheffer (2009), Donadio et al., (2008) and Martins (2009) strengthen the detection of these changes, based on the assumption that infertility has several causes, including obesity, changes in body weight, where these criteria have a negative effect on the conception process.

Problems related to sleep were identified in the literature, and in order to strengthen these findings, the study by Zaidouni et al. (2019) examined the effect of the nursing consultation on stress in infertile couples before starting treatment, and identified 71.4% of the sample with inadequate sleep.

Within the psychobiological level, sexuality and reproduction was the basic human need with the greatest quantitative representativeness, accounting for 20 indicators. This is due to its direct association with procreation, bringing as empirical indicators: situations of obstetric complications, dysfunctions that involve sexual intercourse, such as dissatisfaction, decreased libido and robotization of the sexual act.

Regarding this last indicator, Batista, Bretones and Almeida (2016) observed in their study several complaints related to the robotization of the sexual act, which was seen as a mandatory procedure, an act that suppressed sexual pleasure with their partner. Farias et

al. (2010) corroborate by stating that sex becomes an obligation and this reflects in the disconnection of pleasure, in addition to enabling the reduction of sexual frequency. And in more critical situations, the couple separates themselves and the sexual act is absent (Marques; Morais, 2018).

Table 3. Distribution of empirical indicators regarding the basic human needs of women undergoing assisted reproduction at the psychobiological level and their frequency according to the integrative literature review.

Psychobiological Needs	Empirical indicators	N	%
Nutrition	<ul style="list-style-type: none"> ✓ Inadequate nutrition ✓ Malnutrition ✓ Increased or decreased appetite ✓ Eating disorder ✓ Postnatal eating disorders ✓ Unhealthy eating ✓ Obesity ✓ Peso anormal 20 < IMC < 25 ✓ Excessive caffeine intake 	9	5,42
Elimination	<ul style="list-style-type: none"> ✓ Nausea ✓ Vomiting ✓ Risk of constipation 	3	1,80
Rest and repose	<ul style="list-style-type: none"> ✓ Problems with sleep ✓ Insomnia ✓ Fatigue 	3	1,80
Physical activity	<ul style="list-style-type: none"> ✓ Intense physical activity and exhaustion ✓ Reduced physical activity ✓ Sedentariness 	3	1,8
Sexuality and reproduction	<ul style="list-style-type: none"> ✓ Increased risk of obstetric complications ✓ Increased risk for miscarriage ✓ Increased risk for placenta previa ✓ Increased risk for gestational diabetes ✓ Increased risk for preeclampsia ✓ Increased risk for endometriosis ✓ Increased risk for polycystic ovary syndrome ✓ Increased risk for placental abruption ✓ Increased risk of preterm birth ✓ Increased risk of fetal chromosomal abnormalities ✓ Increased risk of low birth weight ✓ Increased risk for STIs due to multiple partners in order to become pregnant ✓ Sexually Acquired Pelvic Infections ✓ Sexual dysfunction ✓ Underestimating the importance of sexual intimacy in married life ✓ Sexual dissatisfaction / Less satisfaction in sexual intercourse ✓ Quality of impaired sexual intercourse <ul style="list-style-type: none"> ✓ Decreased libido ✓ Orgasm dysfunction ✓ Robotization of the sexual act 	20	12,04

Physical and environmental security	<ul style="list-style-type: none"> ✓ Coping with violence by the partner or family ✓ Living in polluted areas ✓ Increased smoking ✓ Alcohol intake ✓ Drug use 	5	3,01
Regulation: Cell growth and functional development	<ul style="list-style-type: none"> ✓ Exposure to environmental pollutants causes increased oxidative stress ✓ Increased age results in decreased number of follicles in the ovaries and the quality of oocytes 	2	1,20
Vascular regulation	<ul style="list-style-type: none"> ✓ Hypertension ✓ Swelling/Edema 	2	1,20
Neurological regulation	<ul style="list-style-type: none"> ✓ Mood disorder ✓ Obsessive compulsive disorder ✓ Increased risk of psychiatric disorders related to the threat of abortion 	3	1,80
Hormone regulation	<ul style="list-style-type: none"> ✓ Increased insulin level ✓ Hormonal regulation affected by stress ✓ Use of large amounts of hormone repeatedly 	3	1,80
Sensory perception	<ul style="list-style-type: none"> ✓ Colic ✓ Muscle tension ✓ Back pain 	3	1,80
Therapy and prevention	<ul style="list-style-type: none"> ✓ Non-acceptance of the treatment method ✓ Delay in treatment ✓ Low adherence to treatment ✓ Withdrawal from treatment over time ✓ Refusal to perform more invasive tests ✓ Successive treatment failures ✓ Uncertain duration and procedure of treatment ✓ Painful process of self-administration of oral, enteral, parenteral medication <ul style="list-style-type: none"> ✓ Need for psychological intervention ✓ Increased Requirement for Proof of Pregnancy: More USG Exams and Check-Up ✓ Increased risk of cesarean delivery <ul style="list-style-type: none"> ✓ Risk of hysterectomy ✓ Increased risk of cancer due to increased exposure to hormone therapy <ul style="list-style-type: none"> ✓ Problems with ovulation induction <ul style="list-style-type: none"> ✓ Ovulatory changes ✓ Treatment overload ✓ Music therapy ineffective in reducing anxiety ✓ Inadequate professional care at times of abortion or puerperium 	18	10,84
TOTAL		74	44,58

Source: Prepared by the authors.

The data presented in Table 4 below correspond to the empirical indicators of altered BHN related to the psychosocial level. Indicators referring to the needs of: recreation and leisure, space and creativity were not identified in the scientific literature.

Starting with the psychosocial level, which contains the manifestations through instincts such as: living in community, establishing communication, emotions that permeate the individual, among others (Marques et al., 2008). Regarding the empirical indicators related to psychosocial needs, the following were not addressed: recreation and leisure, space and creativity.

As the first category of needs present at the psychosocial level, Communication establishes a close relationship with the next category, Gregaria. This is due to the intimate relationship between the obstacles in communication, which directly reflects on the distance within the relationship.

This is what Marques and Morais (2018) find, when they report that attempts to fail in treatment impacted the inhibition of the spouses, which generated a lack of dialogue between the couple.

The most representative category of the study was emotional safety, as it included 40 empirical indicators, which reinforces the relevance of alluding to items associated with the emotional status of women undergoing assisted reproduction treatments. It is important to highlight that among the indicators detected in the literature there was an explicit plurality of feelings.

In the context of infertility, these women are faced with a situation of need, where it may not lead to a physical affection, however it weakens the psychic health of the couple and even of the entire family (Mesquita et al., 2016). Thus, in view of this complexity that permeates infertility, the need for interventions in the emotional sphere is notorious (Farinati; Rigoni, Muller, 2006).

Table 4. Distribution of empirical indicators regarding the basic human needs of women undergoing assisted reproduction at the psychosocial level and their frequency according to the integrative literature review.

Psychosocial Needs	Empirical indicators	N	%
Communication	✓ Difficulty in communication	1	0,60
Gregarious	<ul style="list-style-type: none"> ✓ Community and family pressure in the face of infertility <ul style="list-style-type: none"> ✓ Bullying parental ✓ Social isolation ✓ Social exclusion, especially of family and spouse <ul style="list-style-type: none"> ✓ Family neglect ✓ Reduction of social and spousal support <ul style="list-style-type: none"> ✓ Deterioration of family relationships ✓ Avoids contact with family and friends with young children <ul style="list-style-type: none"> ✓ Negative social interactions ✓ Damaged marital relationship 	10	6,02

Emotional security	<ul style="list-style-type: none"> ✓ Not believing in the success of the treatment <ul style="list-style-type: none"> ✓ Cultural stigmatization ✓ Depression ✓ Feeling of loss ✓ Suffering ✓ Risk of emotional maladjustment ✓ Emotional pain caused by childlessness <ul style="list-style-type: none"> ✓ Anger ✓ Resentment ✓ Hostile behavior ✓ Sadness ✓ Jealousy ✓ Emotional exhaustion ✓ Prejudice overload ✓ Stress related to marital and personal life <ul style="list-style-type: none"> ✓ Feeling of guilt ✓ Anxiety ✓ Anxiety about the survival of the fetus <ul style="list-style-type: none"> ✓ Anguish ✓ Negative emotions ✓ Grieving process ✓ Despair ✓ Fear of continuing pregnancy ✓ Fear of losing the pregnancy ✓ Fear of miscarriage ✓ Concern associated with decreased fetal movements <ul style="list-style-type: none"> ✓ Fear of divorce ✓ Ectopic Pregnancy Measure ✓ Fear of child malformation ✓ Fear of ineffective treatment ✓ Fear of complications in treatment / Fear about the impact of AR treatment on your health / Fear of continuing treatment due to the high hormonal load <ul style="list-style-type: none"> ✓ Concerns about medical procedures ✓ Avoids expressing emotions ✓ Frustration when treatment fails ✓ Insecurity ✓ Insecurity of having a child gestated in another womb <ul style="list-style-type: none"> ✓ Helplessness ✓ Hopelessness ✓ Distrust ✓ Feeling under constant psychological pressure 	40	24,09
Love, acceptance	<ul style="list-style-type: none"> ✓ Improper coping, avoids the truth about infertility ✓ Difficulty accepting and adapting to reproductive dysfunction ✓ Need for explicit recognition of previous fertility trajectory ✓ Feeling that your family does not accept these methods ✓ Believing that your friends are bored with your infertility 	5	3,01
Self-esteem, self-confidence, self-respect	<ul style="list-style-type: none"> ✓ Low self-esteem ✓ Personal devaluation 	16	9,63

	<ul style="list-style-type: none"> ✓ See yourself as different from those who get pregnant naturally ✓ Lower self-confidence between the couple <ul style="list-style-type: none"> ✓ Decreased self-confidence ✓ Reduced maternal confidence ✓ Feeling vulnerable ✓ Diminished self-image/Damage to self-perception <ul style="list-style-type: none"> ✓ Body depreciation ✓ Lack of femininity ✓ Feeling unable to share feelings <ul style="list-style-type: none"> ✓ Feeling inadequate ✓ Feeling of embarrassment and shame <ul style="list-style-type: none"> ✓ Feeling of failure ✓ Feeling of demand ✓ Loss of control over the body 		
Freedom and participation	<ul style="list-style-type: none"> ✓ Do not have the opportunity to express your concerns during routine check-ups ✓ Interruption of working life 	2	1,20
Health education and learning	<ul style="list-style-type: none"> ✓ Insufficient knowledge about infertility ✓ Superstitions related to infertility ✓ Lack of knowledge about success rates and cost of treatment 	3	1,80
Self-realization	<ul style="list-style-type: none"> ✓ Marital dissatisfaction ✓ Lower life satisfaction ✓ Poor quality of life during treatment ✓ Decreased well-being ✓ Demotivation 	5	3,01
Ensuring access to technology	<ul style="list-style-type: none"> ✓ Late initiation of treatment due to delay in being referred to a specialist in human reproduction ✓ Inaccessible material resources (medications, exams, among others) ✓ Treatment environment has suffered a notable decrease and interaction with professionals is reduced ✓ Difficulty in accessing the program due to the distance from their homes ✓ Difficulty in continuing treatment due to financial issues to bear the high costs of medication and procedures ✓ Limitation of access, due to the advanced age of the woman and her marital status 	6	3,61
TOTAL		88	53,01

Source: Prepared by the authors.

Encompassing the last level of the NHB, the following table explains the 4 HEIs of altered psychospiritual needs.

Within the psychospiritual needs, there is only one representative, Religiosity and spirituality, in which 4 empirical indicators were found in the literature: little religiosity, spiritual confrontation, feeling of divine punishment and feeling of divine injustice.

Thus, people who face serious illnesses have a decrease in stress through the relationship with the spiritual. Likewise, there are hypotheses that the connection with the divine positively influences both the psychological and physical aspects of patients undergoing assisted reproduction treatment, which achieve better results (Braga et al., 2018).

Table 5. Distribution of empirical indicators regarding the basic human needs of women submitted to assisted reproduction at the psychospiritual level and their frequency according to the integrative literature review.

Psychospiritual Needs	Empirical indicators	N	%
Religiosity and spirituality	<ul style="list-style-type: none"> ✓ Little religiosity ✓ Spiritual confrontation ✓ Feeling of divine punishment for mistakes made previously ✓ Sense of divine injustice 	4	2,40

Source: Prepared by the authors.

To emphasize the conceptualization of empirical indicators, Denser (2003) states that in contexts within the normal range or in conditions within the expected range, the indicators expose the changes, and it is from their detection that one can act in the prevention of problems. Thus, this study identified inadequacies of the basic human needs of the public that undergoes assisted reproduction treatments so that it was possible to act emphatically on these points and minimize the damage caused.

CONCLUSION

In contexts within normality or in conditions within expectations, empirical indicators expose the changes, and it is from their detection that one can act to prevent problems. Thus, this study identified inadequacies of the basic human needs of the public that undergoes assisted reproduction treatments so that it was possible to act emphatically on these points and minimize the damage caused.

The study aimed to identify the indicators of women submitted to assisted reproduction in order to structure a software for the systematization of Nursing Care in the light of the Theory of Basic Human Needs, as the scientific knowledge generated by nursing throughout its practice provides a demand for organization, as a result of the progress in the construction and arrangement of conceptual models. Thus, the study had its objectives achieved.

Therefore, the identification of empirical indicators of women undergoing assisted reproduction will support a sovereign care practice, with theoretical foundation and support

through scientific evidence. Thus, the fruits generated by this study will work as a framework for the creation of a software intended for use by the professional team and, more specifically, by nursing, aiming to ensure care that permeates all levels of Basic Human Needs and is concerned with executing the entire Nursing Process.

REFERENCES

1. ANVISA. (2018). 11º Relatório do Sistema Nacional de Produção de Embriões (SISEMBRIO).
2. Batista, L. A. T., Bretones, W. H. D., & Almeida, R. J. (2016). O impacto da infertilidade: narrativas de mulheres com sucessivas negativas pelo tratamento de reprodução assistida. *Reprod. Clim*, 31(3), 121–127.
3. Boivin, J., Bunting, I., Collins, J. A., & Nygren, K. G. (2007). International estimates of infertility prevalence and treatment-seeking: potential need and demand for infertility medical care. *Human Reproduction*, 22(6), 1506-1512.
4. Braga, D. P. A. F., Melamed, R. M. M., Setti, A. S., Zanetti, B. F., Figueira, R. C. S., Assumpção, A., & Borges, J. (2018). Role of religion, spirituality, and faith in assisted reproduction. *Journal of Psychosomatic Obstetrics & Gynecology*, 40(3), 195-201.
5. Brasil. Ministério da Saúde. (2005). Portaria nº 426/GM em 22 de março de 2005. Política Nacional de Atenção Integral em Reprodução Humana Assistida e dá outras providências.
6. CFM. Conselho Federal de Medicina. (2017). Resolução nº 2.168/2017. Dispõe sobre as normas éticas para a utilização das técnicas de reprodução assistida. *Diário Oficial da União, Seção 1, Brasília - DF, p. 73*. Disponível em: <https://sistemas.cfm.org.br/normas/visualizar/resolucoes/BR/2017/2168>.
7. Coelho, M. L. F., et al. (2017). Mídias digitais como auxiliares no processo criativo em design: análise de uso do aplicativo Farbe. *Revista Brasileira de Design da Informação*, 17(1), 106-122.
8. Cook, R. J., & Dickens, B. M. (2014). Reducing stigma in reproductive health. *International Journal of Gynaecology and Obstetrics*, 125(1), 89-92.
9. Daniluk, J. C., & Koert, E. (2013). The other side of fertility coin: A comparison of childless men's and women's knowledge of fertility and assisted reproductive technology. *Fertility and Sterility*, 99(3), 839-846.
10. Denser, C. P. A. C. (2003). Indicadores: instrumento para a prática de enfermagem com qualidade. In *Book AMT. Enfermagem de excelência: da visão à ação* (pp. 123-145). Rio de Janeiro: Editora Guanabara Koogan.
11. Donadio, N. F., et al. (2008). Gestaç o heterot pica: possibilidade diagn stica ap s fertilizaç o in vitro. A prop sito de um caso. *Revista Brasileira de Ginecologia e Obstetr cia*, 30(9), 466–469.
12. Fawcett, J. (2013). Thoughts about conceptual models and measurement validity. *Nursing Science Quarterly*, 26(2), 189-191.

13. Farinati, D. M., Rigoni, M. S., & Müller, M. C. (2006). Infertilidade: um novo campo da psicologia da saúde. *Estudos de Psicologia*, 23(4), 433-439.
14. Frank, A. P. A., Rossi, C. A., Vitorino, T. S., Azambuja, P., & Penna, M. L. F. (2015). Comparative study of regulations of assisted reproduction and abortion laws between Brazil, Germany, Colombia, and France. *Reprodução e Climatério*, 30(2), 77-82.
15. Garcias, T. R., & Cubas, M. R. (2012). Diagnósticos, intervenções e resultados de enfermagem: subsídios para sistematização da prática profissional. Rio de Janeiro: Elsevier.
16. Horta, W. A. (2015). *Processo de enfermagem* (2ª ed.). Rio de Janeiro: Guanabara Koogan.
17. Izzo, C. R., et al. (2015). Reprodução humana: estado atual. *Revista da Associação Médica Brasileira*, 61(6), São Paulo.
18. Machin, R., & Couto, M. T. (2014). "Fazendo a escolha certa": tecnologias reprodutivas, práticas lésbicas e uso de bancos de sêmen. *Physis: Revista de Saúde Coletiva*, 24(4), 1255-1274.
19. Marques, P. P., & Morais, N. A. (2018). La experiencia de parejas infértiles diante de intentos infructuosos de reproducción assistida. *Avances en Psicología Latinoamericana*, 36(2), 299-314.
20. Marques, D. K. A., et al. (2008). Análise da teoria das necessidades humanas básicas de Horta. *Revista de Enfermagem UFPE On-line*, 2(4), 410-416.
21. Martins, M. A. (2009). *Clínica médica* (2ª ed.). São Paulo: Editora Manole, 501-506.
22. Mendonça, C. R., Arruda, J. T., & Amaral, W. N. (2014). Função sexual de mulheres submetidas à reprodução assistida. *Revista Brasileira de Ginecologia e Obstetrícia*, 36(11), Rio de Janeiro.
23. Mesquita, M. O., Trevilato, G. C., Schons, M. S., Riquinho, D. L., Záchia, S., Passos, E. P. (2016). Acesso e permanência das usuárias do Sistema Único de Saúde no Programa de Reprodução Assistida de um hospital universitário, com ênfase na variável raça/cor. *Reprod. Clim*, 31(3), 151-158.
24. Neto, V. L. S., et al. (2019). Diagnósticos da CIPE® de pessoas vivendo com AIDS e Indicadores Empíricos. *Revista Brasileira de Enfermagem*, 72(5).
25. OMS. Organização Mundial de Saúde. (2011). *Mulheres e saúde: evidências de hoje, agenda de amanhã*.
26. Practice Committee of the American Society for Reproductive Medicine. (2013). Definitions of infertility and recurrent pregnancy loss: A committee opinion. *Fertility and Sterility*, 99(1), 63.

27. Polit, D., Beck, C. T., & Hungler, B. P. (2019). Fundamentos de pesquisa em enfermagem: avaliação de evidências para a prática da enfermagem (10ª ed.). Porto Alegre: Editora Artmed.
28. Queiroz, A. B. A., Mohamed, R. P. S., Moura, M. A. V., Souza, I. E. O., Carvalho, M. C. M. P., & Vieira, B. D. G. (2020). Nursing work in assisted human reproduction: between technology and humanization. *Revista Brasileira de Enfermagem*, 73(3).
29. Soares, C. B., Hoga, L. A. K., Peduzzi, M., Sangaleti, C., Yonekura, T., & Delage, Silva, D. R. A. (2014). Revisão integrativa: conceitos e métodos utilizados na enfermagem. *Revista Esc Enferm USP*, 48(2), 335-345.
30. Sommerville, I. (2019). Engenharia de software (10ª ed.). São Paulo: Pearson Education.
31. Yev, I., Harun_or_Rashid, M., Rustamov, O., Zakirova, N., Kasuya, H., & Sakamoto, J. (2012). Social correlates of female infertility in Uzbekistan. *Nagoya Journal of Medical Science*, 74(3-4), 273–283.
32. Zaidouni, A., Ouasmani, F., Benbella, A., Kasouati, J., & Bezaad, R. (2019). The effect of nursing consultation based on Orem's theory of self-care and Bandura's concept on infertility stress. *Journal of Human Reproductive Sciences*, 12(3), 247-254.